

# Psychosis

Australasian Psychiatry  
2014, Vol 22(3) 299  
© The Royal Australian and  
New Zealand College of Psychiatrists 2014  
Reprints and permissions:  
sagepub.co.uk/journalsPermissions.nav  
DOI: 10.1177/1039856214533279  
apy.sagepub.com



1. Psychosis with a later age of onset.

Options:

- A. Adler
- B. Bleuler
- C. Charcot
- D. Jaspers
- E. Jung
- F. Korsakoff
- G. Kraepelin
- H. Roth
- I. Schneider

For each of the following descriptions choose the most appropriate option.

- I. Described a “paraphrenia” group of patients characterised by more affective expression and a relatively benign illness course.
- II. Described “late-onset schizophrenia” as being similar to early-onset schizophrenia, but with an age of onset greater than 40 years.

2. What percentage of all incident cases of major mental illness appear before the age of 25 years?

- A. 1%
- B. 5%
- C. 25%
- D. 50%
- E. 75%

3. When considering referring a patient with schizophrenia for support in finding employment, which of the following would you expect to find in an employment agency with good fidelity to Individual Placement and Support principles?

- A. Clear exclusion criteria
- B. Early discussion regarding the potential loss of pension money
- C. Good integration with mental health services
- D. Gradual transitions to the workplace
- E. Supporting patients in the employment setting for at least three months

### Answers:

- 1. I. G)  
II. B)
- 2. E)
- 3. C)