

Suicide prevention

Australasian Psychiatry 2014, Vol 22(2) 200 © The Royal Australian and New Zealand College of Psychiatrists 2014 Reprints and permissions: sagepub.co.uk/journalsPermissions.nav DOI: 10.1177/1039856214528124 apy.sagepub.com



The following MCQs test knowledge presented in the papers and Editorial relating to suicide in this issue of *Australasian Psychiatry*:

- 1. Mental health literacy is best defined as:
 - A. An awareness of evidence–based practices in mental health
 - B. Education given to patients regarding their disorders
 - C. Public knowledge about mental disorders
 - D. The level of education that mental health workers have received
 - E. The reading ability mental health consumers
- 2. Theories about suicide. Options:
 - A. Anomic suicide
 - B. Durkheim's sociological theory
 - C. Imitative suicides
 - D. Joiner's theory
 - E. Menninger's theory
 - F. "Psychache"
 - G. Psychoanalytic theory
 - H. Savage's egotistical theory
 - I. Stream analogy framework
 - J. The "broken promise effect"

For each of the following descriptions choose the most appropriate explanatory theory.

- i. The assessment of patients' beliefs about their relationships is paramount
- ii. Higher rates of suicide occur on days that are traditionally seen as enjoyable

- iii. The impulse to suicide results from a homicidal drive
- 3. In relation to suicide, an understanding of the concept of "malignant alienation" can help clinicians prevent suicide by ensuring that they?
 - A. Ask suicidal patients to call a 24-hour crisis number
 - B. Attempt to address social isolation in suicidal patients
 - C. Attempt to teach the suicidal patient problem solving skills
 - D. Do not join the suicidal patient's relatives and friends in making disparaging remarks
 - E. Join a peer review group in order to feel supported when treating suicidal patients
- 4. Many measures often used to reduce inpatient suicide have not been validated scientifically. Which of the following measures have been studied and shown to be effective?
 - A. Locking psychiatric units
 - B. "One to one" nursing
 - C. Removing ligature points
 - D. Restricting visiting hours
 - E. Searching patient's items

Answers: 1. C) 2. i. D) ii. J) iii. G) 3. D) 4. C)